

General Information for Hand Surgery Patients

Please read the following information carefully. You have just undergone surgery and there is a dressing and/or splint placed on your surgical site to reduce motion and improve your comfort post-operatively.

Information for after your surgery...

1. **Medication:** Over-the-counter medication can make you more comfortable after your surgery. Tylenol, Extra-strength Tylenol, or Ibuprofen (Advil), can be taken as it is prescribed. During the first 2-3 days, do not wait to feel any discomfort, and plan to take the medication regularly.
2. Keep your hand elevated as much as possible following your surgery. Your hand should be at a level above your heart. When you sleep at night, you can place it on several pillows. This helps decrease the swelling and improve your level of comfort following surgery.
3. If the dressing feels too tight, you can loosen the outer tensor bandage and re-wrap it lightly.
4. You may apply ice on your hand for 15 minutes every few hours during the first few days following surgery. Be sure to not get the splint or dressing wet in doing so.
5. **Dressing Care:** _____

6. You will be unable to perform any heavy lifting or strenuous exercise for a period of time following your surgery. The duration of this will be reviewed during your first follow-up visit.
7. Please call our office today or tomorrow to make a follow-up appointment for _____

8. After you leave the hospital, there are a few things to watch for during your healing process. If you see any of the following, please call our office, or St. Paul's hospital at 604-682-2344 and ask to speak to the plastic surgeon on call:
- Leg swelling
 - Shortness of breath
 - Fever and chills
 - Redness around your incisions
 - Increase pain in your hand

If you have any questions during your post-operative course, please do not hesitate to contact Dr. Seal, or our office.